

	Articulated molts Ichnofossils	Articulated+disarticulated												Limited taxonomic control					<i>Morania</i> *													
		<i>Burgessochaeta</i>	<i>Canadaspis</i>	<i>Canadia</i>	<i>Chancelloria</i>	<i>Eiffelia</i>	<i>Elrathina</i>	<i>Haplophrentis</i>	<i>Liangshanella</i>	<i>Marrella</i>	<i>Micromitra</i>	<i>Olenoides</i>	<i>Selkirkia</i>	<i>Sidneyia</i>	<i>Waptia</i>	<i>Wiwaxia</i>	<i>Mackenzia</i>	<i>Crenorhabdotus</i>		<i>Chaetognath</i>	<i>Hemichordates</i>	<i>Holothurians</i>	<i>Pikaia</i>	<i>Odontogriphus</i>								
BA120																																47%
BA0																																44%
BA-110																																8%
BA-120																																45%
BA-130																																14%
BA-150																																27%
BA-170																																39%
BA-210																																29%
BA-235																																28%
BA-245																																46%
BA-250																																13%
BA-260																																17%
BA-265																																7%
BA-310																																5%
BA-320																																4%
BA-350																																0%
BA-360																																2%
BA-370																																16%
BA-380																																94%
BA-400																																90%
BA-420																																6%
BA-430																																0%
BA-445																																17%
BA-455																																1%
BA-465																																0%
BA-480																																1%

*Ratio presence of *Morania* per field number per bedding assemblage