doi: 10.2110/sedred.2010.2 Volume 8, No. 2, June 2010


ecord

A publication of SEPM Society for Sedimentary Geology with the Sedimentary Geology Division of GSA

The



INSIDE: THE GEOCHEMISTRY OF CAVE CALCITE DEPOSITS AS A RECORD OF PAST CLIMATE PLUS: COUNCIL'S COMMENTS – The Evolving Relevancy of SEPM SEPM RESEARCH CONFERENCE AND WORKSHOP -Paleosols and Surface System Analogs




SEPM BOOKSTORE

Special Publication #94

Application of Modern Stratigraphic Techniques: Theory and Case Histories

Edited by: Kenneth T. Ratcliffe and Brian A. Zaitlin

Much has been written and debated about the various methodologies applied to modern stratigraphic analysis and the ever increasing complexity of terminologies. However, there exist numerous stratigraphic techniques that are reliant upon precise, quantitative, reproducable data, rather than qualitative interpretative stratigraphic methodologies. Such stratigraphic techniques are applied in an entirely pragmatic non-biased manner within the petroleum industry to provide enhanced stratigraphic understanding of petroleum systems. The petroleum industry is a key driver behind the development of new stratigraphic techniques and a major provider of new stratigraphic data, which has resulted in several of these new techniques having been developed as a requirement to the industry. Furthermore, because techniques, such as isotope chemostratigraphy, elemental chemostratigraphy, magnetic susceptibility stratigraphy, numerical biostratigraphy and heavy mineral stratigraphy are based around precise, quantified and reproducible analytical data, they provide an independent means to test the more interpretative stratigraphic methodologies. This volume attempts an overview of stratigraphic methodologies, but largely focuses on data-generative stratigraphic techniques such as chemostratigraphy, magnetic susceptibility stratigraphy, numerical biostratigraphy and heavy mineral stratigraphy. Where appropriate, each paper discusses data generation methods including sample preparation and analytical methods as well outlining data interpretation methods. This is followed by case histories that demonstarte how those data are used to resolve stratigraphic problems, commonly using material derived from petroleum basins around the World.



Catalog #40094 • SEPM Member Price: \$80.00



Now available in CD ROM SEPM Special Publications #1 - #85, Short Course Notes, Core Workshops and Field Trip Guidebooks

SEPM Member Price: \$20 per CD Order from the SEPM Bookstore.


Concepts in Sedimentology and Paleontology #9 Sequence Stratigraphy of Siliciclastic Systems – The ExxonMobil Methodology

Edited by: Vitor Abreu, Jack E. Neal, Kevin M. Bohacs and James L. Kalbas

The stratigraphic concept of a depositional sequence was introduced to the scientific literature by Exxon Production Research Company (EPRco) in the late 70s, building on the shoulders of giants like Chamberlain, Sloss and Wheeler. Since then, several papers compared and contrasted the original Exxon (and later, ExxonMobil) sequence-stratigraphic school with other approaches to subdivide the geologic record, as well as, debating the ExxonMobil model validity and impact on the community. At its core, the ExxonMobil "model" is really a stratigraphic interpretation method, which was never explicitly documented in the literature. The objective of this book is to present the ExxonMobil sequence stratigraphic method in its current form in an attempt to clarify its usage and application in diverse geologic data and depositional environments. This publication is the result of more than 3 decades of sequence stratigraphy research and application at EPRco and at the ExxonMobil Upstream Research Company (URC). The objective is to emphasize the most important aspects of Sequence Stratigraphy – a method to guide geologic interpretation of stratigraphic data (seismic profiles, well-logs, cores and outcrops) across scales (from local to regional and global) and depositional environments (from continental to deep marine). This book in a 11" x 17" format is designed to be easily used for teaching or self-learning experiences.

Catalog #85009 • SEPM Member Prices: Hardbound \$105.00, Softbound \$60.00





Cover image: Photo of wide stalagmites and flowstone developed in Niedzwiedzia Cave, Kletno, Poland. Paleo-water levels are indicated by calcite deposition along the old water surface; this deposition is clearly visible on the broad stalagmite in the center of the image.

<u>CONTENTS</u>

- 4 The geochemistry of cave calcite deposits as a record of past climate
- **10** Council's Comments The Evolving Relevancy of SEPM
- **11** Research Conference and Workshop Paleosols and Surface System Analogs

The Sedimentary Record (ISSN 1543-8740) is published quarterly by the Society for Sedimentary Geology with offices at 4111 S. Darlington, Suite 100, Tulsa , OK 74135-6373, USA.

Copyright 2010, Society for Sedimentary Geology. All rights reserved. Opinions presented in this publication do not reflect official positions of the Society.

The Sedimentary Record is provided as part of membership dues to the Society for Sedimentary Geology.

Editors

Ruarri Day-Stirrat ruarri.day-stirrat@beg.utexas.edu Bureau of Economic Geology, University of Texas – Austin Xavier.Janson@beg.utexas.edu Bureau of Economic Geology, University of Texas – Austin Wayne Wright wayne.wright@beg.utexas.edu Bureau of Economic Geology, University of Texas – Austin

SEPM Staff

4111 S. Darlington, Suite 100, Tulsa, OK 74135-6373 Phone (North America): 800-865-9765 Phone (International): 918-610-3361

Dr. Howard Harper, Executive Director hharper@sepm.org Theresa Scott, Associate Director & Business Manager tscott@sepm.org Michele Tomlinson, Publication & Technology Coordinator mtomlinson@sepm.org Janice Curtis, Membership Associate jcurtis@sepm.org Edythe Ellis, Administrative Assistant eellis@sepm.org

SEPM Council

Paul M. (Mitch) Harris, President MitchHarris@Chevron.com **Christopher Fielding**, President-Elect cfielding2@unl.edu Diane Kamola, Secretary-Treasurer kamola@ku.edu Maria Mutti, International Councilor mmutti@geo.uni-potsdam.de Nancy Engelhardt-Moore, Councilor for Paleontology nengelhardt-moore@comcast.net Evan Franseen, Councilor for Sedimentology evanf@kgs.ku.edu Samuel Bentley, Councilor for Research Activities sbentley@mun.ca Paul McCarthy, Co-Editor, JSR mccarthy@gi.alaska.edu Gene Rankey, Co-Editor, JSR grankey@ku.edu Stephen Hasiotis, Co-Editor, PALAIOS hasiotis@ku.edu Edith Taylor, Co-Editor, PALAIOS etaylor@ku.edu Gary Nichols, Co-Editor, Special Publications g.nichols@gl.rhul.ac.uk Brian Ricketts, Co-Editor, Special Publications Brian.ricketts@xtra.co.nz Tim Carr, President, SEPM Foundation tim.carr@mail.wvu.edu

www.sepm.org

The geochemistry of cave calcite deposits as a record of past climate

James U.L. Baldini

Department of Earth Sciences, Durham University, Durham DH1 3LE, United Kingdom james.baldini@durham.ac.uk doi: 10.2110/sedred.2010.2.4

ABSTRACT

Cave calcite deposits, and stalagmites specifically, are proving to be critical recorders of paleoclimate. Stalagmites develop from the slow accumulation of calcite deposited after the degassing of carbon dioxide from percolation water in caves. Stalagmites are therefore theoretically capable of recording the hydrochemistry of every drip of water that impacts their surface, but complications involving hydrology, cave air PCO₂ variability, intra-aquifer processing, and petrography preclude straightforward interpretation. Nevertheless, stalagmite climate proxy records using stable isotopes, particularly of oxygen, are reconstructing the timing, structure, and in some cases magnitude of climate shifts worldwide. Trace element to calcium ratios in stalagmite are a promising new proxy for both independent reconstruction of climate and for supporting oxygen isotope ratios. Extensive monitoring of individual cave sites is required to most accurately interpret stalagmite proxy records.

INTRODUCTION

Speleothems are secondary sedimentary, usually monomineralic, cave deposits that may consist of any mineral and may take on any number of morphologies. Over 250 different minerals have been identified in caves, ranging from common species such as calcite and gypsum to mush less common species, such as vanadate and organic minerals (Hill and Forti 1997). Despite this broad diversity of minerals, speleothems are predominantly composed of only three: calcite, aragonite, and gypsum, and of these the vast majority are composed of calcite. Of the different forms of speleothem (including stalagmites, stalactites, helictites, flowstones, and many more), calcite stalagmites are by far the most researched and provide the vast majority of important climate records (Fig 1) (McDermott 2004). Stalagmites are formed by the gradual accumulation of calcite from a drip emanating from the point where a water flowpath through the aquifer intersects a void space. Groundwater in karst aquifers typically has very high levels of dissolved carbon dioxide and calcium due to contact with the soil derived CO₂ (Murthy et al. 2003) and subsequent limestone dissolution (White 1988). When the drip enters the cave atmosphere, which almost always has a lower CO_2 partial pressure (PCO_2) than the soil zone with which the water has equilibrated with, CO₂ degassing occurs from the drip resulting in calcite deposition. Cave drips therefore typically become oversaturated with respect to calcite immediately upon exposure to the cave atmosphere, and deposit calcite first on

the ceiling of the cave, resulting in a stalactite. After the drip gains enough mass to overcome surface tension (Collister and Mattey 2008), it falls to the cave floor where it deposits calcite until it completes its equilibration with ambient P_{CO_2} . The relatively straightforward morphology of calcite stalagmites (Fig 2), combined with their occurrence worldwide and amenability to radiometric dating, means that they are excellent multi-proxy archives of climate, particularly in parts of the world where other easily-dated climate archives (e.g., ice cores, deep sea sediments cores) are scarce or nonexistent (Cruz et al. 2006; Hodge et al. 2008; McDermott et al. 2001).

This paper reviews how stalagmite geochemistry records climate, and discusses some case studies to demonstrate the ability of these sediments to provide both long-term and very highly resolved records. In addition, research aiming to better constrain how the climate signal is altered as groundwater passes through the soil, epikarst, and bedrock, until its discharge point in the cave is summarized. Finally, some future research problems, challenges, and applications of the technique will be introduced.

KARST HYDROLOGY

Karst landscapes are characterized by an abundance of caves, sinkholes (dolines), and lack of surface drainage. Karst aquifers are often problematic for hydrological studies because the basic rules of hydrogeology as defined by Darcy's Law sometime do not apply. Groundwater flow is often non-Darcian in karst aquifers; that is, it is more comparable to open channel flow and therefore often adheres more closely to the Manning Equation than Darcy's Law. Because stalagmites grow in the unsaturated zone of the aquifer, much of the relevant groundwater flow is vertical and may be affected by local anisotropic features such as bedding planes, fractures, and dissolutional features. However, abundant intragranular permeability may also exist, as well as smaller fractures that may store water for long time intervals. Some estimates suggest that the majority of water stored in the unsaturated zone of karst aquifers is stored in this 'diffuse' permeability reservoirs (Atkinson 1977). This leads to a situation where stalagmites may be fed by drips with vastly different water histories: i) water that have resided in the aquifer for decades or longer, *ii*) drips that have only recently percolated through the aquifer, or *iii*) any combination of the two. Recharge to growing stalagmites is therefore generally composed of diffuse and fracture-fed components and identifying the relative importance of both of these is critical for understanding how climate is conveyed from the atmosphere to the stalagmite.




Figure 1. Calcite cave deposits in Niedzwiedzia Cave, Kletno, Poland. The stalagmites accumulate from the floor upwards and the stalactites from the roof downwards.

Recent research has demonstrated that different stalagmites record climate with differing resolution (Baldini et al. 2006a; Cruz et al. 2005) depending on the hydrochemical characteristics of the drip feeding the sample. Different stalagmites, even within the same cave, can record different aspects of the climate signal; some samples may retain a long-term averaged climate signal while others may record individual rain events. Overly responsive drips result in stalagmites yielding very 'noisy' records, can be subject to unpredictable flow rerouting, and may be seasonally undersaturated with respect to calcite, and should therefore be avoided for anything except very specialized applications. Conversely, diffuse drips result in stalagmites with an extremely 'averaged' climate signal that only encodes a small percentage of total rainfall isotope variability, and, while useful for determining longer-term climate, are unsuitable for reconstructing seasonal-scale or event-scale climate (Baldini et al. 2006b; McDermott et al. 2006). Morphological characteristics of stalagmites combined with drip monitoring using automated drip rate loggers (Mattey et al. 2008) and dripwater

chemical analyses (Baldini et al. 2006a; Tooth and Fairchild 2003) help identify samples that may yield the climate signal most suitable for specific project goals. For example, several drip sites in Crag Cave, Castleisland, Ireland, were evaluated and compared to rainfall amounts to better understand the different hydrologies that could occur within one cave (Fig 3). Diffuse, responsive, and overly responsive drip sites were identified, and it was determined that only stalagmites underneath the diffuse and responsive drips would be suitable for a climate reconstruction study (Baldini et al., *unpublished data*).

CAVE ATMOSPHERE CONTROLS ON CALCITE DEPOSITION

Although hydrological controls on stalagmite growth and climate signal emplacement have been very well researched recently, a less well quantified variable critical to stalagmite growth is the differential between soil and cave air PCO_2 . Because research (Baldini et al. 2008; Banner et al. 2007) has demonstrated that cave air P_{CO_2} greatly affects the rate of calcite deposition, characterization of cave air PCO2 variability is required to aid in interpretation of any climate records produced. There are numerous ways that the CO₂ may enter cave systems, but in most temperate limestone caves the CO₂ is derived from the soil zone, either as direct percolation of gaseous CO2 through fractures or indirectly dissolved in drip water. Without any ventilation, the cave would eventually reach the same P_{CO_2} as the soil zone and no degassing, and no calcite deposition, would occur. Cave ventilation, controlled by changes in temperature, air density, barometric pressure, or wind direction, is known to modulate cave air PCO2. Because drip equilibration with ambient PCO2 results in the deposition of calcite, elevated cave air P_{CO_2} results in less calcite deposition than low cave air PCO2. For example, New St. Michael's Cave in Gibraltar ventilates in the summer (Mattey et al. 2008), so, assuming other growth determining variables remain constant, more calcite deposition during the summer months would be expected. This would skew annually averaged climate reconstructions towards summer recharge, and thus impacting the net climate signal (Baldini et al. 2008). To assess the seasonality of calcite deposition, loggers are often used to record cave air PCO2, thus facilitating the interpretation of any proxy records generated. New studies are attempting to measure all growth determining variables while simultaneously collecting stalagmite calcite to test the concepts outlined above (Fig 4).

The **Sedimentary** Record

OXYGEN ISOTOPE RATIOS

The majority of paleoclimatic records using stalagmites are based on the ratio of stable isotopes within the calcite, particularly the ratio of $^{\rm 18}{\rm O}$ to $^{\rm 16}{\rm O}$, or the $\delta^{\rm 18}{\rm O}.$ Stalagmite δ^{18} O is derived from the drip water, which itself is rainwater processed within the karst aquifer and soil. Rainwater δ^{18} O is largely affected by temperature, the rainout amount, and the δ^{18} O of water from the moisture source region (itself often linked to climate), so stalagmites record changes in these parameters. Stalagmite growth occurs gradually from calcite precipitated directly from cave drip water; stalagmite geochemistry is therefore dependent on drip water hydrochemistry which in turn reflects rainwater chemistry and climate. Because drip water δ^{18} O reflects rainfall δ^{18} O (McDermott 2004), stalagmites preserve a record of past variability in rainfall δ^{18} O. The initial focus of studies using $\delta^{\scriptscriptstyle 18}O$ in stalagmites was paleotemperature reconstruction; if calcite and drip water $\delta^{_{18}}O$



Figure 2. A cross-sectional view of stalagmite BA-99-2 from Ballynamintra Cave, Dungarvan, Ireland. The calcite is deposited as lens that drape over each previous lens. Typically analytical traverses run along the central growth axis of the stalagmite, as this is the location that experiences the least amount of kinetic fractionation and has the highest deposition rate and therefore the highest temporal resolution achievable using microanalytical techniques. The sample is approximately 30cm long.




Figure 3. Automatically logged drip rates for three drip sites in Crag Cave, Ireland. A) An example of a diffuse flow drip. Note the static drip rate and lack of response to rainfall events. B) An intermediate 'responsive' drip site. Note how large rainfall events result in a rapid increase in drip rate followed by a more gradual decrease. A diffuse flow component is also present, demonstrated by the asymptotic behaviour of the recessional limb of the storm hydrographs. C) An example of an overly responsive drip site. The water responds to nearly all rain events within hours and is often undersaturated with respect to calcite.

are known, the temperature that the calcite was deposited at can be calculated, because the water-calcite fractionation factor for oxygen isotopes is temperature dependent (Craig 1965). Although this undoubtedly does affect calcite δ^{18} O, external climate exerts a much stronger influence on the drip water $\delta^{_{18}}O$ than cave air temperature, therefore stalagmite $\delta^{_{18}}O$ records are now generally interpreted as reflecting shifts in rainwater δ^{18} O. Some of the most important climate records are derived from regions that experience extremely variable amounts or types of rainfall. Because rainwater derived from different sources, occurring at different times of year, or that is impacted by the 'amount effect' (Rozanski et al. 1993), can have vastly different δ^{18} O values, shifts in any of these parameters can be manifest in stalagmite records. Southeast Asia

experiences a strongly monsoonal climate, and thus stalagmites from the region record variations in monsoonal strength through time (Wang et al. 2001). Monsoonal rainfall is characterized by extremely low δ^{18} O values compared to non-monsoon rainfall, so changing proportions of monsoonal recharge compared to non-monsoonal rainfall results in fluctuations in groundwater recharge δ^{18} O. Increased monsoonal strength causes lower δ^{18} O values of groundwater and stalagmite calcite. Combined with the excellent chronological control afforded by U-series techniques, this permits extraordinarily long records of monsoonal variability to be constructed for this region, which can then be compared with incoming solar radiation to achieve a better understating of the timing of major shifts in the Earth's climate on both

orbital and suborbital timescales (Cheng et al. 2009; Cheng et al. 2006; Henderson 2006).

Another good example of the applicability of stalagmites to reconstruct climate comes from the North Atlantic, where tropical cyclone (TC) rainfall is characterized by having anomalously low δ^{18} O values. Variability in TC activity is dependent on a vast number of natural and anthropogenic factors, so that definitively ascribing causality to global warming (or naturally occurring variability) is problematic. Recent statistical research attributes increasing TC intensity to increasing sea surface temperatures (SSTs) caused by global warming (Emanuel 2005; Hoyos et al. 2006; Webster et al. 2005), but uncertainty regarding data quality and the effects of natural decadal-scale climate oscillations have greatly complicated attribution (Landsea 2005; Pielke 2005). Furthermore, two recent climate model studies actually predict reductions in Atlantic TC frequency assuming predicted 21st Century global warming estimates (Knutson et al. 2008). No doubt exists that Atlantic TC activity has increased since 1995, but the 20th century record suggests that similarly elevated levels also existed from 1920-1969 (Goldenberg et al. 2001). Much of the contentiousness results from the brevity and questionable reliability of the available records. Accurate records of total Atlantic TC activity only extend back to the beginning of routine aircraft reconnaissance (1944-present), while somewhat reliable estimates extend back to the 1900s (Goldenberg et al. 2001), greatly reducing the statistical significance of correlations between Atlantic TC activity and climatic parameters. Rainfall in humid, low-latitude regions is characterised by elevated δ^{18} O values, typically ranging from 0 to -5‰ SMOW (Dansgaard 1964). Intense rainfall produced by TCs deviates from this trend, and is characterised by extremely negative δ^{18} O values (typically -8 to -14 ‰ SMOW) (Lawrence et al. 1998) and provides a substantial amount of groundwater recharge in areas affected by these storms. Simple calculations demonstrate that one TC alone can lower integrated annual groundwater δ^{18} O values by 1.5‰ in a localized area and by 0.5‰ over an area several hundreds of kilometres across (Fig 5). An even greater isotopic anomaly would result in a stalagmite if the stalagmite is analysed at subannual resolution. Because stalagmite $\delta^{18}O$ is correlated directly with unsaturated zone groundwater δ^{18} O, stalagmites preserve a record of past variability in total rainfall amount derived from these seasonal, anomalously low δ^{18} O rainfall events. A recent




Figure 4. An SEM image of secondary calcite (A and B) deposited on an Iceland Spar (C) substrate placed underneath the drip CC-Bil (see figure 3). Well-developed columnar crystals are apparent (B). The calcite grew over the course of six months. All growth determining variables were measured simultaneously, so our understanding of calcite accumulation rates can be tested (Sherwin and Baldini, unpublished data). Scale bar division are 50 µ.

study from Belize used a very high-resolution $\delta^{_{18}}$ O record from Actun Tunichil Muknal Cave in Belize to demonstrate that not only was the record a reflection of the El Niño-Southern Oscillation, but that brief isotopic excursions in the record coincided with hurricanes passing near the site (Frappier et al. 2007). The record only stretches back to 1977, but it clearly demonstrates the power and usefulness of this technique; the record consists of approximately one data point every month and detects every hurricane during the interval of calcite growth. Other research is trying to extend this record. For example, a \$1.6 million European Research Councilfunded project (HURRICANE) based at the University of Durham (UK) aims to reconstruct TC activity throughout the Caribbean region using stalagmites. The principal goal of the HURRICANE Project is to reconstruct TC activity extending back at least 500 years, greatly lengthening existing records and helping to ascertain whether the current observed increases in TC activity are the result of anthropogenic global warming or part of a natural cycle.

TRACE ELEMENTS

Whereas oxygen isotope ratios are used to reconstruct changes in the seasonal moisture balance, temperature, and moisture source δ^{18} O; trace element ratios of stalagmite calcite are used to reconstruct paleohydrology, paleorainfall, and potentially paleo-bioproductivity of soils. Despite much promise and research, the controls on the trace element geochemistry of stalagmites remain comewhat enigmatic. Stalagmite calcite Mg and Sr concentrations may record paleo-recharge conditions due to a combination of incongruent dissolution, selective leaching, and residence time mechanisms (Fairchild et al. 2000; Huang et al. 2001; McDonald et al. 2004; McMillan et al. 2005; Roberts et al. 1998; Treble et al. 2003), and these trace elements have historically been the focus of most of the research output. Studies suggest that for karst aquifers without dolomite, groundwater residence time in the aquifer is a critical control on Mg and Sr concentrations in cave drip waters (Fairchild et al. 2000). A multiyear drip water study of an Australian cave system suggested that drip water Mg/Ca and Sr/Ca at this site increase dramatically during droughts (McDonald et al. 2004). This increase may reflect more CO2 degassing in an increased volume of void spaces in the aquifer,

The **Sedimentary** Record

subsequently leading to increased calcite precipitation above the stalagmite. Because of low partition coefficients in calcite of Mg and Sr, this process reduces drip water [Ca2+] but does not impact drip water [Mg2+] and [Sr2+], resulting in elevated ratios. This 'prior calcite precipitation' (PCP) mechanism has been demonstrated to be an important influence on Mg and Sr concentrations in drip water in other studies as wel' (Baldini et al. 2006a; Fairchild et al. 2000; Tooth and Fairchild 2003), and our recent unpublished work establishes the link between hydrology, cave air PCO₂, and PCP. Long-term monitoring studies of a wide range of drip types show that PCP amounts vary according to site hydrological characteristics (Baldini et al. 2006a; Tooth and Fairchild 2003), and research in the Edwards aquifer of central Texas also suggests that increased groundwater residence times and fluid-rock interactions result in elevated drip water Mg/Ca and Sr/Ca, and that these trace elements are therefore good indicators of reduced recharge at these sites (Musgrove and Banner 2004). However, research has predominantly focused on using the trace elements Mg and Sr as paleoclimate proxies, either independently or to corroborate δ^{18} O based paleoclimatic interpretations.




Figure 5. An example of the effects of a burricane on regional isotope hydrology. (A) The geographic extent of Hurricane Rita at the moment of landfall September 24, 2005 (adapted from NOAA satellite data). The large size of tropical cyclones means that broad areas are affected by low δ^{18} O rainfall. (B) The estimated effect of Hurricane Rita on groundwater δ^{18} O values. The values shown are the per mil difference between hypothetical groundwater not affected by any burricanes (calculated assuming all contributions to groundwater came from non-tropical rainfall) and hypothetical groundwater affected just by Hurricane Rita, which lowered integrated annual groundwater δ^{18} O by approximately 1.5 % SMOW near the Texas/Louisiana border and by 0.5 ‰ over a much broader area. Stalagmite δ^{18} O values would be lowered by the same amount (Mickler et al. 2006). The values were calculated using measured Hurricane Rita rainfall amounts, mean annual rainfall for the area shown, and approximate δ^{18} O of burricane (Lawrence et al. 1998) and non-burricane rainfall (Dansgaard 1964).




Figure 6. A very high-resolution strontium concentration transect through calcite that may have been deposited during the '8.2 ka' cold dry event from a stalagmite obtained from Crag Cave, Ireland (adapted from Baldini etal., 2002). Lower strontium values are representative of wetter conditions, and elevated values of drier conditions. Seven annual cycles are represented by the trace element data, which refines the uranium-series chronology used to define the x-axis that suggests just five years of growth. The lower growth rate suggested by the strontium data compared to the lower spatial resolution uranium series chronology is consistent with cold, dry conditions prevalent over this short time interval, which would have discouraged calcite deposition.

Microanalytical techniques have successfully resolved annual cycles in stalagmite trace element records (Baldini et al. 2002; Fairchild et al. 2001; Treble et al. 2003) at scales as low as 1 Ìm, providing a potentially powerful method of building annual-scale chronologies and reconstructing paleoseasonality (Fig 6). Borsato et al. (2007) used micro-X-ray fluorescence spectrometry (Ì-XRF) to study Pb, Zn, Fe, Sr, P, Br, Cu, and Y annual cycles at the micron-scale in a stalagmite from Grotta di Ernesto (northern Italy) and assessed the strength of the annual signal for these elements. A Secondary Ionization Mass Spectrometry (SIMS) of several different speleothems identified annual scale cycles in Mg, Sr, Ba, H, F, Na, and P (Fairchild et al. 2001), suggesting that annual trace element cycles are not uncommon. Of these, Fairchild et al. (2001) suggest that P may have the greatest potential as a paleoclimate proxy due to its role as a nutrient element. This was supported by a study of a modern Australian stalagmite that matched trace element cycles to the instrumental record and found that P, U, and Mg were the most reliable paleohydrological indicators at that site (Treble et al. 2003). However, the low soil retention capacity for P at that site may have increased the effectiveness of P as a paleohydrological proxy (Treble et al. 2003).

Despite the potential importance of P as a paleoclimate indicator in stalagmite calcite,

interpretations have largely depended on inferential relationships with other proxies and calcite petrography; very little information on P concentrations or seasonality in cave drip waters exists. Drip water geochemical work has suggested that several elements, not just P, may be transported bound to colloidal material (Borsato et al. 2007). However, no long term results on the P variability of a cave drip exist. Clearly the interest in applying trace element cycles to reconstruct paleoclimate stems largely from the very high resolution achievable using modern microanalytical

techniques such as SIMS, µ-XRF, excimer laser ablation inductively coupled plasma mass spectrometry (ELA-ICPMS), and micromilling. However, it is also apparent that drip water sampling studies have not matched the weeklyto monthly- scale temporal resolution possible through microanalytical analyses of speleothem samples, largely because of the logistical difficulties involved in collecting samples daily or even weekly. Automated drip rate loggers have produced very high resolution records of cave drip discharge rates (Collister and Mattey 2008; Hu et al. 2008; McDonald and Drysdale 2007; Sondag et al. 2003) and conductivity (Genty and Deflandre 1998), but long-term drip water trace element monitoring studies have remained at the monthly-scale. Consequently, high-resolution drip water trace element data are critical missing information that would help clarify the links between stalagmite trace element concentrations, climate, aquifer-related processes (e.g., residence time), and crystallographic mechanisms (e.g., calcite growth rate).

CONCLUSION AND FUTURE CHALLENGES

Stalagmites are proving to be exceptionally valuable repositories of climate information for low- to mid-latitude terrestrial settings where few other long-term, high-resolution, and accurately dateable climate proxies are found. Although the initial promise of deriving absolute paleotemperatures from stalagmite calcite is currently unachievable, stable isotope ratios of oxygen and trace element-to-calcium ratios are proving to be excellent indicators of shifts in moisture availability, bioproductivity, and the strength of isotopically anomalous precipitation events such as hurricanes and monsoons. The excellent chronological controls achievable by using U-230 Th dating techniques, combined with high precision multi-collector ICP-MS analyses, (McDermott 2004) means that, although stalagmite climate proxies may not typically permit quantification of the amplitude of environment change, climatic shifts can be identified and the timing, duration, and structure of climatic events can be reconstructed. Oxygen isotope ratios combined with complementary trace element data provide a powerful tool for unraveling past climates. Ongoing research using stalagmites is reconstructing climate on a variety of timescales, from the high-profile research on the Asian Monsoon that has repercussions for global paleoclimate on millennial to orbital timescales (Cheng et al. 2009; Johnson et al. 2006), to new research such as the HURRICANE Project that is attempting to reconstruct more recent paleoclimate variability on event scales. Despite the success of stalagmites in reconstructing global paleoclimate, a number of challenges still exist. It may yet be possible to use stalagmite calcite as a paleothermometer, and new techniques using clumped isotopes (Meckler et al. 2009) and fluid inclusions (Vonhof et al. 2006) are promising. Cave monitoring studies are critical for understanding how the climate signal is conveyed through the aquifer to the stalagmite calcite. The processes controlling the incorporation of many trace elements in stalagmite calcite are still poorly understood and these records are best interpreted after considerable monitoring and site characterization. It is becoming clear that every cave has a unique set of controls that modulate how the climate signal is recorded in stalagmite calcite, and a set protocol on how to best delineate these controls has yet to be established. Despite these challenges, stalagmites are providing key climate proxy records that are greatly improving our understanding of past climate fluctuations, which has important consequences for predicting future change.

ACKNOWLEDGMENT

Dr. Lisa Baldini is thanked for providing images and useful discussion. Catherine Sherwin is thanked for providing SEM images of calcite growth. The landowners of Crag Cave (Ireland), Ballynamintra Cave (Ireland), and

Niedzwiedzia Cave (Poland) are thanked for providing continued access to the cave sites. The editors are also thanked for their efficient handling of the manuscript.

REFERENCES

- ATKINSON, T.C., 1977, Diffuse flow and conduit flow in limestone terrain in the Mendip Hills, Somerset (Great Britain): Journal of Hydrology, v. 35, p. 93-110.
- BALDINI, J.U.L., MCDERMOTT, F., and FAIRCHILD, I.J., 2002, Structure of the 8200-year cold event revealed by a speleothem trace element record: Science, v. 296, p. 2203-2206.
- BALDINI, J.U.L., MCDERMOTT, F., and FAIRCHILD, I.J., 2006a, Spatial variability in cave drip water hydrochemistry: Implications for stalagmite paleoclimate records: Chemical Geology, v. 235, p. 390-404.
- BALDINI, J.U.L., MCDERMOTT, F., HOFFMANN, D.L., RICHARDS, D.A., and CLIPSON, N., 2008, Very high-frequency and seasonal cave atmosphere PCO₂ variability: Implications for stalagmite growth and oxygen isotope-based paleoclimate records: Earth and Planetary Science Letters, v. 272, p. 118-129.
- BALDINI, L.M., MCDERMOTT, F., and BALDINI, J.U.L., 2006b, Detecting NAO-mode variability in high-resolution speleothem isotope records: Geochimica et Cosmochimica Acta, v. 70, p. 1.
- BANNER, J.L., GUILFOYLE, A., JAMES, E.W., STERN, L.A., and MUSGROVE, M., 2007, Seasonal variations in modern speleothem calcite growth in Central Texas, USA: Journal of Sedimentary Research, v. 77, p. 615-622.
- BORSATO, A., FRISIA, S., FAIRCHILD, I.J., SOMOGYI, A., and SUSINI, J., 2007, Trace element distribution in annual stalagmite laminae mapped by micrometer-resolution X-ray fluorescence: implications for incorporation of environmentally significant species: Geochimica et Cosmochimica Acta, v. 71, p. 1494-1512.
- CHENG, H., EDWARDS, R.L., BROECKER, W.S., DENTON, G.H., KONG, X.G., WANG, Y.J., ZHANG, R., and WANG, X.F., 2009, Ice Age Terminations: Science, v. 326, p. 248-252.
- CHENG, H., EDWARDS, R.L., WAN, Y.J., KO, X.G., MING, Y.F., KELLY, M.J., WANG, X.F., GALLUP, C.D., and LIU, W.G., 2006, A penultimate glacial monsoon record from Hulu Cave and two-phase glacial terminations: Geology, v. 34, p. 217-220.

COLLISTER, C., and MATTEY, D., 2008, Controls on water drop volume at speleothem drip sites: An experimental study: Journal of Hydrology, v. 358, p. 259-267.

CRAIG, H., 1965, The measurement of oxygen isotope palaeotemperatures, in Tongiorgi, E., ed., Stable Isotopes in Oceanographic Studies and Palaeotemperatures: Pisa, Consigleo Nazionale della Ricerche Laboratorio di Geologia Nucleare, p. 3-24.

- CRUZ, F.W., BURNS, S.J., KARMANN, I., SHARP, W.D., VUILLE, M., and FERRARI, J.A., 2006, A stalagmite record of changes in atmospheric circulation and soil processes in the Brazilian subtropics during the Late Pleistocene: Quaternary Science Reviews, v. 25, p. 2749-2761.
- CRUZ, F.W., KARMANN, I., VIANA, O., BURNS, S.J., FERRARI, J.A., VUILLE, M., SIAL, A.N., and MOREIRA, M.Z., 2005, Stable isotope study of cave percolation waters in subtropical Brazil: Implications for paleoclimate inferences from speleothems: Chemical Geology, v. 220, p. 245-262.
- DANSGAARD, W., 1964, Stable isotopes in precipitation: Tellus, v. 16, p. 436-468.
- EMANUEL, K., 2005, Increasing destructiveness of tropical cyclones over the past 30 years: Nature, v. 436, p. 686-688.
- FAIRCHILD, I.J., BAKER, A., BORSATO, A., FRISIA, S., HINTON, R.W., MCDERMOTT, F., and TOOTH, A.F., 2001, Annual to sub-annual resolution of multiple traceelement trends in speleothems: Journal of the Geological Society of London, v. 158, p. 831-841.

FAIRCHILD, I.J., BORSATO, A., TOOTH, A.F., FRISIA, S., HAWKESWORTH, C.J., HUANG, Y., MCDERMOTT, F., and SPIRO, B., 2000, Controls on trace element (Sr-Mg) compositions of carbonate cave waters: implications for speleothem climatic records: Chemical Geology, v. 166, p. 255-269.

FRAPPIER, A.B., SAHAGIAN, D., CARPENTER, S.J., GONZALEZ, L.A., and FRAPPIER, B.R., 2007, Stalagmite stable isotope record of recent tropical cyclone events: Geology, v. 35, p. 111-114.

GENTY, D., and DEFLANDRE, G., 1998, Drip flow variations under a stalactite of the Père Noël cave (Belgium). Evidence of seasonal variations and air pressure constraints: Journal of Hydrology, v. 211, p. 208-232.

GOLDENBERG, S.B., LANDSEA, C.W., MESTAS-NUNEZ, A.M., and GRAY, W.M., 2001, The recent increase in Atlantic hurricane activity: Causes and implications: Science, v. 293, p. 474-479.

HENDERSON, G.M., 2006, Climate - Caving in to new chronologies: Science, v. 313, p. 620-622.

HILL, C., and FORTI, P., 1997, Cave Minerals of the World: Huntsville: National Speleological Society.

HODGE, E.J., RICHARDS, D.A., SMART, P.L., ANDREO, B., HOFFMANN, D.L., MATTEY, D.P., and GONZALEZ-RAMON, A., 2008, Effective precipitation in southern Spain (similar to 266 to 46 ka) based on a speleothem stable carbon isotope record: Quaternary Research, v. 69, p. 447-457.

HOYOS, C.D., AGUDELO, P.A., WEBSTER, P.J., and CURRY, J.A., 2006, Deconvolution of the factors contributing to the increase in global hurricane intensity: Science, v. 312, p. 94-97.

HU, C.Y., HENDERSON, G.M., HUANG, J.H., CHEN, Z.H., and JOHNSON, K.R., 2008, Report of a three-year monitoring programme at Heshang Cave, Central China: International Journal of Speleology, v. 37, p. 143-151.

- HUANG, Y., FAIRCHILD, I.J., BORSATO, A., FRISIA, S., CASSIDY, N.J., MCDERMOTT, F., and HAWKESWORTH, C.J., 2001, Seasonal variations in Sr, Mg and P in modern speleothems (Grotta di Ernesto, Italy): Chemical Geology, v. 175, p. 429-448.
- JOHNSON, K.R., LYNN INGRAM, B., SHARP, W.D., and ZHANG, P., 2006, East Asian summer monsoon variability during Marine Isotope Stage 5 based on speleothem ¥18O records from Wanxiang Cave, central China: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 236, p. 5-19.

KNUTSON, T.R., SIRUTIS, J.J., GARNER, S.T., VECCHI, G.A., and HELD, I.M., 2008, Simulated reduction in Atlantic hurricane frequency under twenty-firstcentury warming conditions: Nature Geoscience, v. 1, p. 359-364.

LANDSEA, C.W., 2005, Hurricanes and global warming: Nature, v. 438, p. E11-E13.

LAWRENCE, J.R., GEDZELMAN, S.D., ZHANG, X.P., and ARNOLD, R., 1998, Stable isotope ratios of rain and vapor in 1995 hurricanes: Journal of Geophysical Research-Atmospheres, v. 103, p. 11381-11400.

MATTEY, D., LOWRY, D., DUFFET, J., FISHER, R., HODGE, E., and FRISIA, S., 2008, A 53 year seasonally resolved oxygen and carbon isotope record from a modern Gibraltar speleothem: reconstructed drip water and relationship to local precipitation: Earth and Planetary Science Letters, v. doi: 10.1016/j.epsl.2008.01.051.

MCDERMOTT, F., 2004, Palaeo-climate reconstruction from stable isotope variations in speleothems: a review: Quaternary Science Reviews, v. 23, p. 901-918.

MCDERMOTT, F., BALDINI, L.M., and BALDINI, J.U.L., 2006, Forward modelling of speleothem delta O-18: A critical assessment of their use as high-resolution climate proxies: Geochimica Et Cosmochimica Acta, v. 70, p. A408-a408.

MCDERMOTT, F., MATTEY, D.P., and HAWKESWORTH, C., 2001, Centennial-scale Holocene climate variability revealed by a high-resolution speleothem ¥18O record from SW Ireland: Science, v. 294, p. 1328-1331. MCDONALD, J., and DRYSDALE, R., 2007, Hydrology of cave drip waters at varying bedrock depths from a karst system in southeastern Australia: Hydrological Processes, v. 21, p. 1737-1748.

MCDONALD, J., DRYSDALE, R., and HILL, D., 2004, The 2002-2003 El Nino recorded in Australian cave drip waters: Implications for reconstructing rainfall histories using stalagmites: Geophysical Research Letters, v. 31, p. 1-4.

MCMILLAN, E.A., FAIRCHILD, I.J., FRISIA, S., BORSATO, A., and MCDERMOTT, F., 2005, Annual trace element cycles in calcite-aragonite speleothems: evidence of drought in the western Mediterranean 1200-1100 yr BP: Journal of Quaternary Science, v. 20, p. 423-433.

MECKLER, A.N., ADKINS, J.F., EILER, J.M., and COBB, K.M., 2009, Constraints from clumped isotope analyses of a stalagmite on maximum tropical temperature change through the late Pleistocene: Geochimica et Cosmochimica Acta, v. 73, p. A863-A863.

- MICKLER, P.J., STERN, L.A., and BANNER, J.L., 2006, Large kinetic isotope effects in modern speleothems: Geological Society of America Bulletin, v. 118, p. 65-81.
- GOOBELA SOCKY OF FINICAL DIRECT A DRIVEN WITH, P. 109-01. MURTHY, R., GRIFFIN, K.L., ZARNOCH, S.J., DOUGHERTY, P.M., WATSON, B., VAN HAREN, J., PATTERSON, R.L., and MAHATO, T., 2003, Carbon dioxide efflux from a 550 m3 soil across a range of soil temperatures: Forest Ecology and Management, v. 178, p. 311-327.
- MUSGROVE, M., and BANNER, J.L., 2004, Controls on the spatial and temporal variability of vadose dripwater geochemistry: Edwards Aquifer, central Texas: Geochimica et Cosmochimica Acta, v. 68, p. 1007-1020.

PIELKE, A.R., 2005, Are there trends in hurricane destruction?: Nature, v. 438, p. E11.

- ROBERTS, M.S., SMART, P.L., and BAKER, A., 1998, Annual trace element variations in a Holocene speleothem: Earth and Planetary Science Letters, v. 154, p. 237-246.
- ROZANSKI, K., ARAGUAS-ARAGUAS, L., and GONFIANTINI, R., 1993, Isotopic patterns in modern global precipitation, *in Swart*, P.K., Lohmann, K.C., McKenzie, J., and Savin, S., eds., Climate Change in Continental Isotopic Records: Washington, D.C., American Geophysical Union, p. 1-36.
- SONDAG, F., VAN RUYMBEKE, M., SOUBIES, F., SANTOS, R., SOMERHAUSEN, A., SEIDEL, A., and BOGGIANI, P., 2003, Monitoring present day climatic conditions in tropical caves using an Environmental Data Acquisition System (EDAS): Journal of Hydrology, v. 273, p. 103-118.
- TOOTH, A.F., and FAIRCHILD, I.J., 2003, Soil and karst aquifer hydrological controls on the geochemical evolution of speleothem-forming drip waters, Crag Cave, southwest Ireland: Journal of Hydrology, v. 273, p. 51-68.
- TREBLE, P., SHELLEY, J.M.G., and CHAPPELL, J., 2003, Comparison of high resolution sub-annual records of trace elements in a modern (1911-1992) speleothem with instrumental climate data from southwest Australia: Earth and Planetary Science Letters, v. 216, p. 141-153.
- VONHOF, H.B., VAN BREUKELEN, M.R., POSTMA, O., ROWE, P.J., ATKINSON, T.C., and KROON, D., 2006, A continuous-flow crushing device for on-line delta H-2 analysis of fluid inclusion water in speleothems: Rapid Communications in Mass Spectrometry, v. 20, p. 2553-2558.
- WANG, Y.J., CHENG, H., EDWARDS, R.L., AN, Z.S., WU, J.Y., SHEN, C.-C., and DORALE, J.A., 2001, A high-resolution absolute-dated late Pleistocene monsoon record from Hulu Cave, China: Science, v. 294, p. 2345-2348.
- WEBSTER, P.J., HOLLAND, G.J., CURRY, J.A., and CHANG, H.R., 2005, Changes in tropical cyclone number, duration, and intensity in a warming environment: Science, v. 309, p. 1844-1846.
- WHITE, W.B., 1988, Geomorphology and Hydrology of Karst Terrains: New York, Oxford University Press.

Accepted April 2010

COUNCIL'S COMMENTS

The Evolving Relevancy of SEPM

Many of us joined SEPM before the technological revolution placed access to its two flagship journals - Journal of Sedimentary Research and PALAIOS - at the fingertips of almost anyone with access to a computer keyboard. In those days, membership in SEPM meant receiving our own copies of these journals at our desks, and made remaining up-to-date on current research trends much more convenient than a trip to the library. SEPM membership has always helped to support these journals and to ensure dissemination of the highest quality science in sedimentary geology and paleontology. But SEPM has always been much more than just its journals.

SEPM continues to publish a comprehensive, high quality Special Publications series that highlights thematic works of particular contemporary relevance. All publications are now available on CDs and also as downloadable files from the SEPM website (www.sepmonline.org). In addition, SEPM has always focused on continuing education by sponsoring technical sessions, short courses, workshops and field trips at the AAPG and GSA annual meetings. These SEPM activities provide a variety of forums for members to disseminate the results of their work as well as providing accessibility to new ideas, techniques and directions in sedimentary geology and paleontology. SEPM also provides discounts to its members to attend meetings and to purchase publications, and for students, an opportunity to secure funding for their research.

As the digital age permeates further into our everyday professional lives, the existence of a community of geoscientists with common interests is perhaps more relevant than ever. SEPM is the largest and most premiere society emphasizing sedimentary geology and paleontology in North America and it also includes a significant international membership. The international footprint of SEPM is increasing, and will continue to do so, through the expansion of Global Ambassadors and a stronger presence at key international meetings. SEPM provides a key connection to this community of geoscientists whose activities are relevant to basic scientific inquiry as well as energy and environmental concerns. This community of geoscientists working in academia, industry and government provides critically important networking opportunities that foster the exchange of research ideas and potential employment opportunities.

SEPM is unique among geosciences societies, incorporating both academic and applied, cutting-edge science in sedimentary geology and paleontology. The knowledge our members generate is more relevant to society than ever as we continue to explore for increasingly scarce resources, manage environmental issues and try to understand and adapt to global climate change. SEPM has a unique opportunity to play a leadership role in increasing public awareness and shaping public policy on these issues. In fact, in many respects the future and direction of our Society will be driven by them.

From SEPM Council Members Paul (Mitch) Harris, Paul McCarthy, Gene Rankey



SEPM Society for Sedimentary Geology "Bringing the Sedimentary Geology Community Together" www.sepm.org

2011 Medalists

Medals and awards are an important part of the Society's mission. It is with great pride that we announce the 2011 awardees. They will be honored at the 2011 President's Reception and Awards Ceremony, during the SEPM Annual Meeting held in Houston, Texas.

- Twenhofel Medalist: Walter Dean
- Shepard Medalist: Miriam Kastner
- Pettijohn Medalist: David Rubin
- Moore Medalist: Barun Sen Gupta
- James Lee Wilson Award: Richard Twitchett

Honorary Member: Dag Nummedal



SEPM RESEARCH CONFERENCE AND WORKSHOP Paleosols and Surface System Analogs



Conference Leaders: Steve Driese¹, Lee Nordt¹, Stacy Atchley¹, Steve Dworkin¹, Dan Peppe¹, Cindy Stiles², Curtis Monger³ and Eugene Kelley⁴



¹Department of Geology, Baylor University, Waco, TX, Steven_Driese@baylor.edu

²U.S. Dept. of Agriculture, Natural Resources Conservation Service, National Laboratory, Lincoln, NE

³Plant and Environmental Sciences Dept., New Mexico State University, Las Cruces, NM

⁴Dept. of Soil and Crop Sciences, Colorado State University, Fort Collins, CO

Registration Costs TBA

Introduction and Purpose:

Paleosols are increasingly recognized by researchers and utilized mainly to make inferences regarding past climates (precipitation and temperature), landscapes, and paleoatmospheric chemistry, including pCO₂ and pO₂. Recognition and interpretation of pre-Quaternary paleosols requires the use of multidisciplinary approaches that involve integration of field morphological, microscopic, and geochemical data, as well as an appreciation for evolutionary changes in terrestrial biota and flora that have occurred over time. There is also the increasing use of USDA Soil Taxonomy for paleosols preserved in the rock record. SEPM, the Baylor Terrestrial Paleoclimatology Research Group, and NSF are sponsoring a combined field and research conference September 21-26, 2010, centered at the Petrified Forest National Park near Holbrook, AZ that addresses these six key central questions:

Question 1: What is the preservation potential of different types of paleosols in the geologic record?

Question 2: What is the fidelity of time and event preservation recorded in paleosols?

Question 3: Where do paleosols "fit" in sequence-stratigraphy?

Question 4: Is the current paleoclimate "toolkit" comprised of field morphological and geochemical proxies as advanced as it can likely become?

Question 5: To what extent can standard USDA soil physical and chemical characterization techniques be used for establishing colloidal properties of paleosols?

Question 6: Is paleosol research adequately represented within the US National Research Council-National Science Foundation "Critical Zone" Initiative?

Other Potential Topical Issues:

Taxonomic systems - Paleobotany/Trace fossils/Ecosystems - Regional paleopedology - Paleosol/modeling correlations -

Primary Deliverables Resulting From Conference:


•Based upon a combination of talks, poster sessions and field trips addressing the central questions proposed previously, develop a summary report "white paper" on the status of paleosol research and on future research directions and funding opportunities.

Develop a subset of the submitted contributions into an SEPM Special Publication Volume.



Dates and Location

September 21-26, 2010 at Petrified Forest National Park, near Holbrook in northeastern Arizona, USA



General Research Conference Trip Dates and Schedule:

September 21st (Tuesday): Group arrives Albuquerque International Airport, NM; PM in motel near Airport (TBA).

September 22nd (Wednesday): Group drives 3.5 hours to Holbrook, AZ to check into motel (TBA); afternoon visiting Petrified Forest Visitor Center, various scenic and tourist stops; dinner in Holbrook (at registrant's cost); 8:00 PM PEFO overview provided by Bill Parker, NPS.

September 23rd (Thursday): Plenary Sessions (AM) and Poster Sessions (PM); Barbecue and Mixer at PEFO Picnic Area.

September 24th (Friday): Field Trip (all day) with time to roam at lunch stop.

September 25th (Saturday): Plenary Sessions (AM) and Poster Sessions (PM), with wrap-up white paper draft completed; End-of-conference banquet at Mesa Sports Bar and Grill, Holbrook, AZ.

September 26th (Sunday): Drop registrants at Albuquerque International Airport, NM for return flights, or those with rental cars or private vehicles depart.

Anticipated high temperatures during conference: upper 80's to low 90's (Fahrenheit) with lows in 30's to 40's and very low humidity, with little chance for rain.

SEPM Book Series Online

www.sepmonline.org Introductory Free Access Period

SEPM Books Series are now available online, chapter by chapter, in a similar format to the SEPM journals, JSR and PALAIOS. The main page is located at **www.sepmonline.org**, where access to the four major books series, Special Publication; Concepts in Sedimentology and Paleontology; Short Course Notes and Core Workshop Notes is available.

- Features include Chapter by chapter PDFs, mouse-over abstract previews, free frontmatter and backmatter files, free reference lists with links.
- These archives contain all books in each series from volume 1 through books published in 2005
- Newer books will be added each year
- During the introductory period from April until June, access was free to everyone.
- After June 1st, access is by subscription or by pay-per-view only.
- Yearly subscription rates run from June 1 to May 31 and for 2010-2011 are:
 - Libraries, Institutions and Companies: \$1,200 single site access to all four book series per year
 - SEPM Members (individuals): \$400 access to all four book series per year
 - Subcriptions to individual book series is also available
- Resolution of the online files varies as files may have had to be down sampled allow for efficient download times
- Each book is also available on CD at full resolution. see SEPM Book Catalogue and website (www.sepm.org)
- Contact SEPM with any questions or to subscribe

